

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/198240>

Please be advised that this information was generated on 2019-06-02 and may be subject to change.

EXTENDED GENOME REPORT

Open Access



The draft genome sequence of “*Nitrospira lenta*” strain BS10, a nitrite oxidizing bacterium isolated from activated sludge

Dimitra Sakoula¹ , Boris Nowka², Eva Spieck², Holger Daims³ and Sebastian Lücker^{1*}

Abstract

The genus *Nitrospira* is considered to be the most widespread and abundant group of nitrite-oxidizing bacteria in many natural and man-made ecosystems. However, the ecophysiological versatility within this phylogenetic group remains highly understudied, mainly due to the lack of pure cultures and genomic data. To further expand our understanding of this biotechnologically important genus, we analyzed the high quality draft genome of “*Nitrospira lenta*” strain BS10, a sublineage II *Nitrospira* that was isolated from a municipal wastewater treatment plant in Hamburg, Germany. The genome of “*N. lenta*” has a size of 3,756,190 bp and contains 3968 genomic objects, of which 3907 are predicted protein-coding sequences. Thorough genome annotation allowed the reconstruction of the “*N. lenta*” core metabolism for energy conservation and carbon fixation. Comparative analyses indicated that most metabolic features are shared with *N. moscoviensis* and “*N. defluvii*”, despite their ecological niche differentiation and phylogenetic distance. In conclusion, the genome of “*N. lenta*” provides important insights into the genomic diversity of the genus *Nitrospira* and provides a foundation for future comparative genomic studies that will generate a better understanding of the nitrification process.

Keywords: “*Nitrospira lenta*”, Nitrite oxidation, *Nitrospira*, Wastewater treatment

Introduction

Nitrification, the two-step oxidation of ammonia to nitrate via nitrite, is a key process of the biogeochemical nitrogen cycle. Nitrite-oxidizing bacteria (NOB) are chemolithoautotrophic microorganisms that catalyze the oxidation of nitrite to nitrate, the second step of the nitrification process. For decades NOB were considered as metabolically restricted microorganisms solely associated with nitrification. However, experimental findings contradict this opinion, indicating a versatile ecophysiology of many NOB [1–4] and highlighting their important role in and possibly outside of the nitrogen cycle [5].

The genus *Nitrospira* is the most diverse known NOB genus and is divided in six different phylogenetic sublineages [6–8]. Members of the genus are ubiquitously present in different natural and engineered ecosystems [5, 9–11]. Despite their high abundance, only eleven

representative species, distributed within the six *Nitrospira* sublineages, have been obtained in enrichment or pure culture so far [7, 8, 12–15]. Sublineage I and II *Nitrospira* are considered to be the dominant NOB in most wastewater treatment plants (WWTPs), playing a key role in the efficient removal of nitrogen via nitrification [6, 16]. Besides their widespread distribution and crucial role, the physiology of *Nitrospira* species is highly understudied, mainly due to the lack of pure cultures and genomic data [3, 17, 18]. The recent identification of complete ammonia-oxidizing (comammox) *Nitrospira* [15, 19] not only redefined the nitrification process, but also further indicated the importance of the genus and emphasized our poor understanding of the metabolic versatility present within this phylogenetic group.

“*Nitrospira lenta*” strain BS10 was isolated from a municipal WWTP [13] and it is the fourth isolate belonging to the sublineage II *Nitrospira*, besides *N. moscoviensis* [20], “*N. japonica*” [14], and the comammox organism “*N. inopinata*” [12]. Thus, insights into the “*N. lenta*” genome will shed light onto the genomic flexibility and metabolic

* Correspondence: s.luecker@science.ru.nl

¹Department of Microbiology, IWW, Radboud University, Heyendaalseweg 135, 6525 AJ Nijmegen, Netherlands

Full list of author information is available at the end of the article



diversity of the genus *Nitrospira* and aid further comparative studies between *Nitrospira* species.

Organism information

Classification and features

“*N. lenta*” strain BS10 is a Gram negative, aerobic NOB isolated from activated sludge of a municipal WWTP in Hamburg, Germany (basic properties are summarized in Table 1) [13]. Based on 16S rRNA gene-based phylogenetic analysis, “*N. lenta*” is affiliated with *Nitrospira* sublineage II but is only distantly related to the sublineage II type strain, *N. moscoviensis* (Fig. 1).

Table 1 Classification and general features of “*Nitrospira lenta*” strain BS10 [34]

MIGS ID	Property	Term	Evidence code ^a
	Classification	Domain Bacteria	TAS [35]
		Phylum <i>Nitrospirae</i>	TAS [36]
		Class “ <i>Nitrospira</i> ”	TAS [36]
		Order “ <i>Nitrospirales</i> ”	TAS [36]
		Family “ <i>Nitrospiraceae</i> ”	TAS [36]
		Genus <i>Nitrospira</i>	TAS [32]
		Species <i>Nitrospira lenta</i>	TAS [13]
		Strain: BS10	TAS [13]
	Gram stain	Negative	TAS [13]
	Cell shape	Spiral-shaped rods	TAS [13]
	Motility	Non-motile ^b	TAS [13]
	Sporulation	Not reported	NAS
	Temperature range	10–32 °C	TAS [13]
	Optimum temperature	28 °C	TAS [13]
	pH range; Optimum	7.4–8.0; 7.4–7.6	TAS [13]
	Carbon source	Carbon dioxide	TAS [13]
MIGS-6	Habitat	Wastewater treatment plant	TAS [13]
MIGS-6.3	Salinity	0.5% w/v	TAS [13]
MIGS-22	Oxygen requirement	Aerobic	TAS [13]
MIGS-15	Biotic relationship	Free-living	TAS [13]
MIGS-14	Pathogenicity	Non-pathogen	NAS
MIGS-4	Geographic location	Germany/Hamburg	TAS [13]
MIGS-5	Sample collection	12/12/2006	TAS [13]
MIGS-4.1	Latitude	53° 31′ 8″ N	TAS [13]
MIGS-4.2	Longitude	9° 54′ 53″ E	TAS [13]
MIGS-4.4	Altitude	–	NAS

^aEvidence codes – IDA Inferred from Direct Assay, TAS Traceable Author Statement (i.e., a direct report exists in the literature), NAS Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [37]

^bGenes encoding for a flagellum were identified in “*N. lenta*” BS10 genome

A pure culture of “*N. lenta*” was obtained after applying a combination of standard isolation methods (density gradient centrifugation and serial dilutions) with an optical tweezer system for the sorting of single cells [13]. “*N. lenta*” grows mainly planktonic and forms helical-shaped cells. Cells are 1.0–2.3 µm in length and 0.2–0.3 µm in diameter (Table 1, Fig. 2). As shown by Nowka et al. [13], “*N. lenta*” is able to grow at lower temperatures (10 °C) than *N. moscoviensis*. Interestingly, while “*N. lenta*” is not able to tolerate high concentrations of nitrite, it exhibits a lower affinity for nitrite compared to *N. moscoviensis* and “*N. defluvii*”, which indicates a clear niche differentiation among these *Nitrospira* species [21].

Genome sequencing information

Genome project history

“*N. lenta*” was selected for whole genome sequencing on the basis of its relevance within the nitrogen cycle as well as due to the general lack of genomic information for *Nitrospira* species. Furthermore, because “*N. lenta*” was isolated from activated sludge, its genome was expected to yield insights that allow optimization and stabilization of the nitrification process in wastewater treatment. The draft genome sequence of “*N. lenta*” BS10 was completed on 27/07/2013. The high-quality draft genome is available in the European Nucleotide Archive (ENA) under study accession number PRJEB26290. An overview of the genome sequencing project is given in Table 2.

Growth conditions and genomic DNA extraction

“*N. lenta*” was cultivated as described by Nowka et al. [13] in mineral salt medium amended with 0.02 g L^{−1} NaNO₂[−] as energy source. The cultures were incubated in the dark at 28 °C, with moderate stirring (100–300 rpm). The genomic DNA was extracted following the hexadecyltrimethylammoniumbromide (CTAB) protocol provided by the DOE Joint Genome Institute (JGI, <https://jgi.doe.gov/user-program-info/pmo-overview/protocols-sample-preparation-information/>) as described elsewhere [22].

Genome sequencing and assembly

High-throughput sequencing was performed at GATC Biotech (Constance, Germany) using Roche GS FLX technology. The final draft genome of “*N. lenta*” was obtained using the GS De Novo Assembler (Newbler) and comprised 3.8 Mb on 22 contigs. Genome completeness was evaluated with CheckM [23]. Similarly to the complete genomes of “*N. defluvii*” (98% completeness, 2.3% contamination) and *N. moscoviensis* (96% completeness, 6.6% contamination), the “*N. lenta*” genome was estimated to be 96% complete with 3.2% contamination.





Table 2 Project information


MIGS ID	Property	Term
MIGS 31	Finishing quality	High quality draft
MIGS-28	Libraries used	1
MIGS 29	Sequencing platforms	Roche GS FLX
MIGS 31.2	Fold coverage	40
MIGS 30	Assemblers	GS De Novo Assembler
MIGS 32	Gene calling method	AMIGene [24]
	Locus Tag	NITLEN
	GenBank ID	OUNR00000000.1
	GenBankDate of Release	4 June 2018
	GOLD ID	–
MIGS 13	BioProject	PRJEB26290
	Source Material Identifier	BS10
	Project relevance	Microbiology, Biotechnology

Genome annotation

The draft genome of “*N. lenta*” was annotated using the MicroScope platform [24] as described in detail elsewhere [17]. The automatic annotation was manually checked and curated using the MicroScope Web interface MaGe [25]. The genomic features of “*N. lenta*” were compared to *N. moscoviensis* and “*N. defluvii*”, the type strains of the *Nitrospira* sublineages II and I, respectively, using the OrthoVenn web service [26] for the identification and comparison of orthologous gene groups. Sequence similarities were calculated using an E-value of 1e-05. An inflation value of 1.5 was applied to generate the orthologous clusters.

Genome properties

The “*N. lenta*” draft genome consists of 22 contigs and has a total size of 3,756,190 bp with an overall G + C



content of 57.9% (Fig. 3, Table 3). From a total of 3968 predicted genes, 3907 (98.5%) and 61 (1.5%) are protein and RNA coding sequences, respectively. The genome encodes for 1 complete rRNA operon and 46 tRNAs, with 1 to 5 copies for each tRNA type. Moreover, 66.8% of the predicted genes were assigned into Clusters of Orthologous Groups (COG) functional categories (Table 4).

Insights from the genome sequence

Nitrospira species belonging to sublineages I and II are the most abundant NOB in many environments and play a key role in N-cycling in engineered ecosystems [6, 10]. Recent experimental data indicates a clear niche differentiation between sublineage I and II *Nitrospira* [21, 27]. More specifically, “*N. lenta*”, like other members of sublineage II, exhibits a lower maximum activity [21, 27] and could be outcompeted by sublineage I *Nitrospira* at higher nitrite concentrations [28]. Despite their ecophysiological differences, sublineage I and II *Nitrospira* exhibit substantial genomic similarities. More specifically, “*N. lenta*” shares a core genome including 2223 orthologous protein clusters with *N. moscoviensis* and “*N. defluvi*”. This corresponds to 67.3% of the pan-genome of the *Nitrospira* species included in this analysis (Fig. 4). Moreover, “*N. lenta*” features the lowest number of unique genes (1100, of which 51 are grouped in 18 paralogous protein clusters). Most of these unique genes lack any function prediction (Fig. 4).

Table 3 Genome statistics

Attribute	Value
Genome size (bp)	3,756,190
DNA coding (bp)	3,487,097
DNA G + C (%)	57.9
DNA scaffolds	22
Total genes	3968
Protein coding genes	3907
RNA genes	61
rRNA genes	3
tRNA genes	46
Pseudo genes	0
Genes in internal clusters	–
Genes with function prediction	1990
Genes assigned to COGs	2609
Genes with Pfam domains	3868
Genes with signal peptides ^a	196
Genes with transmembrane helices	965
CRISPR repeats	0


^aOnly signal peptides with a prediction probability greater 70% were taken into consideration

Table 4 Number of genes associated with general COG functional category prediction

Code	Value	%age ^a	Description
J	163	4.17	Translation, ribosomal structure and biogenesis
A	0	0	RNA processing and modification
K	148	3.79	Transcription
L	142	3.64	Replication, recombination and repair
B	1	0.03	Chromatin structure and dynamics
D	48	1.23	Cell cycle control, Cell division, chromosome partitioning
V	44	1.13	Defense mechanisms
T	256	6.55	Signal transduction mechanisms
M	303	7.76	Cell wall/membrane biogenesis
N	114	2.92	Cell motility
U	119	3.05	Intracellular trafficking and secretion
O	151	3.87	Posttranslational modification, protein turnover, chaperones
C	205	5.25	Energy production and conversion
G	133	3.40	Carbohydrate transport and metabolism
E	198	5.07	Amino acid transport and metabolism
F	59	1.51	Nucleotide transport and metabolism
H	136	3.48	Coenzyme transport and metabolism
I	76	1.95	Lipid transport and metabolism
P	156	3.99	Inorganic ion transport and metabolism
Q	77	1.97	Secondary metabolites biosynthesis, transport and catabolism
R	389	9.96	General function prediction only
S	221	5.66	Function unknown
–	1298	33.22	Not in COGs

^aThe total is based on the total number of protein coding genes in the genome

In accordance with its ability to oxidize nitrite to nitrate [13], “*N. lenta*” encodes all proteins required for nitrite oxidation (Fig. 5), for which the key enzyme is a membrane-associated nitrite oxidoreductase (NXR) [29]. This protein complex belongs to the type II dimethyl sulfoxide reductase family of molybdopterin-binding enzymes and consists of three subunits [17, 29]. The “*N. lenta*” genome contains two paralogous copies of *nxrA* and *nxB*, encoding the NXR α and β subunits, respectively, and two copies of *nxC* for the candidate γ subunit. Like all *Nitrospira* genomes analyzed to date, the “*N. lenta*” genome contains *nirK*, encoding the copper-dependent NO-forming nitrite reductase. While the function of NirK in *Nitrospira* is still unclear, a role in dissimilatory nitrite reduction is unlikely as no other genes involved in denitrification were identified in “*N. lenta*” or any other *Nitrospira*. One cannot exclude the possibility that NO plays a regulatory role in *Nitrospira*, for example in the regulation of forward versus reverse



electron flow as proposed for *Nitrobacter* [30]. Moreover, "*N. lenta*" exhibits the genetic capacity for nitrogen assimilation from nitrite as its genome features *nirA* encoding the ferredoxin-dependent nitrite reductase. Interestingly, NirA is conserved in "*N. defluvii*", but not the other genome-sequenced sublineage II *Nitrospira*, which either encode an octaheme nitrite reductase [3, 18], or, in the case of the comammox *Nitrospira*, lack assimilatory nitrite reduction pathways [15, 19, 31]. Interestingly, the "*N. lenta*" BS10 genome also features an *ure* operon encoding a functional urease (UreC), as well as a complete gene set (*urtABCDE*) for a high affinity urea ABC transporter. This implies that "*N. lenta*" is able to hydrolyse urea to ammonium and CO₂, facilitating nitrogen and carbon assimilation from urea and reciprocal feeding between "*N. lenta*" and urease-negative ammonia-oxidizing bacteria [3, 4]. The "*N. lenta*" urease is closely related to the enzyme of *N. moscoviensis*, but significantly differs from known ureases of ammonia-oxidizing bacteria [3].


"*N. lenta*" conserves energy by nitrite oxidation with oxygen as terminal electron acceptor. During nitrite oxidation catalyzed by NXR, two electrons are shuttled (putatively via cytochrome *c*) towards a putative novel *bd*-like terminal oxidase [17]. The proton motive force established through active proton pumping by this novel complex IV and/or the release of scalar protons during nitrite oxidation drives ATP synthesis by a F_oF₁-type ATPase (complex V). The other respiratory complexes

(complexes I to III) will not contribute to energy conservation during lithoautotrophic growth on nitrite, but will operate in reverse to provide reducing equivalents for carbon fixation [17]. Moreover, the complete gene repertoire for the oxidative and reductive tricarboxylic acid (TCA) cycle is present in "*N. lenta*" for pyruvate oxidation via acetyl-CoA and CO₂ fixation, respectively. Moreover, the complete glycolysis/gluconeogenesis and pentose phosphate pathways were identified. The observed presence of glycolysis and the oxidative TCA cycle might indicate that "*N. lenta*" can benefit from a mixotrophic lifestyle in the presence of organic carbon, as has been reported for other *Nitrospira* representatives [6, 16, 32].

Finally, the "*N. lenta*" genome contains various defense mechanisms against heavy metals, antibiotics, and the antibacterial agent acriflavine. "*N. lenta*" encodes a superoxide dismutase for defense mechanisms against oxidative stress, as well as several bacterioferritins, which can participate in oxidative stress resistance mechanisms [17].

Conclusions

Together with *N. moscoviensis* and "*N. japonica*", "*N. lenta*" represents only the third cultured species of canonical nitrite-oxidizing *Nitrospira* from sublineage II. In this study, the genome of "*N. lenta*" was analyzed, demonstrating that "*N. lenta*" shares a significant amount of genomic features with other representatives of the genus. However,



physiological differences observed by Nowka et al. regarding growth conditions and nitrite affinities [13, 21] clearly suggest a niche differentiation between different species. The “*N. lenta*” genome will facilitate a better understanding of the metabolic versatility of the genus *Nitrospira* and will be useful for future comparative studies, especially those with a focus on species obtained from engineered systems.

Abbreviations

NOB: Nitrite-oxidizing bacteria; NXR: Nitrite oxidoreductase; WWTP: Wastewater treatment plant

Acknowledgements

We are grateful to Lianna Poghosyan, Jeroen Frank and Maartje van Kessel for helpful discussions. The LABGeM (CEA/IG/Genoscope & [CNRS UMR8030](#)) and the France Génomique National infrastructure (funded as part of Investissement d'avenir program managed by Agence Nationale pour la Recherche, contract ANR-10-INBS-09) are acknowledged for support within the MicroScope annotation platform.

Funding

This work was supported by the European Research Council (ERC AG 339880 ECOMOM), the Netherlands Organisation for Scientific Research (NWO; VENI grant 863.14.019), the Vienna Science and Technology Fund (WWTF; grant LS09-40), the Austrian Science Fund (FWF; grants P24101-B22 and P25231-B21), and the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) grant SP 667/10-1 + 2).

Authors' contributions

ES, HD, and SL designed the study, DS, BN, and SL performed research, DS and SL analyzed data, DS, ES, HD and SL wrote the paper, all authors reviewed and agreed with the final version of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Microbiology, IWW, Radboud University, Heyendaalseweg 135, 6525 AJ Nijmegen, Netherlands. ²Department of Microbiology & Biotechnology, University of Hamburg, Ohnhorststr. 18, 22609 Hamburg, Germany. ³Division of Microbial Ecology, Department of Microbiology and Ecosystem Science, University of Vienna, Althanstr. 14, 1090 Vienna, Austria.

Received: 12 June 2018 Accepted: 10 November 2018

Published online: 22 November 2018

References

1. Füssel J, Lückers S, Yilmaz P, Nowka B, van Kessel MAHJ, Bourceau P, et al. Adaptability as the key to success for the ubiquitous marine nitrite oxidizer *Nitrospoccus*. *Sci Adv*. 2017;3(11):e1700807.
2. Koch H, Galushko A, Albertsen M, Schintmeister A, Gruber-Dorninger C, Lückers S, et al. Growth of nitrite-oxidizing bacteria by aerobic hydrogen oxidation. *Science*. 2014;345(6200):1052–4.

3. Koch H, Lückner S, Albertsen M, Kitzinger K, Herbold C, Spieck E, et al. Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus *Nitrospira*. *Proc Natl Acad Sci U S A*. 2015;112(36):11371–6.
4. Palatinszky M, Herbold C, Jehmlich N, Pogoda M, Han P, von Bergen M, et al. Cyanate as an energy source for nitrifiers. *Nature*. 2015;524(7563):105–8.
5. Daims H, Lückner S, Wagner M. A new perspective on microbes formerly known as nitrite-oxidizing bacteria. *Trends Microbiol*. 2016;24(9):699–712.
6. Daims H, Nielsen JL, Nielsen PH, Schleifer KH, Wagner M. In situ characterization of *Nitrospira*-like nitrite-oxidizing bacteria active in wastewater treatment plants. *Appl Environ Microbiol*. 2001;67(11):5273–84.
7. Lebedeva EV, Alawi M, Maixner F, Jozsa PG, Daims H, Spieck E. Physiological and phylogenetic characterization of a novel lithoautotrophic nitrite-oxidizing bacterium, '*Candidatus Nitrospira bockiana*'. *Int J Syst Evol Microbiol*. 2008;58(Pt 1):242–50.
8. Lebedeva EV, Off S, Zumbiegel S, Kruse M, Shagzhina A, Lucker S, et al. Isolation and characterization of a moderately thermophilic nitrite-oxidizing bacterium from a geothermal spring. *FEMS Microbiol Ecol*. 2011;75(2):195–204.
9. Freitag TE, Chang L, Clegg CD, Prosser JL. Influence of inorganic nitrogen management regime on the diversity of nitrite-oxidizing bacteria in agricultural grassland soils. *Appl Environ Microbiol*. 2005;71(12):8323–34.
10. Juretschko S, Timmermann G, Schmid M, Schleifer K-H, Pommerening-Röser A, Koops H-P, et al. Combined molecular and conventional analyses of nitrifying bacterium diversity in activated sludge: *Nitrosococcus mobilis* and *Nitrospira*-like bacteria as dominant populations. *Appl Environ Microbiol*. 1998;64(8):3042–51.
11. Lebedeva EV, Alawi M, Fiencke C, Namsaraev B, Bock E, Spieck E. Moderately thermophilic nitrifying bacteria from a hot spring of the Baikal rift zone. *FEMS Microbiol Ecol*. 2005;54(2):297–306.
12. Kits KD, Sedlacek CJ, Lebedeva EV, Han P, Bulaev A, Pjevac P, et al. Kinetic analysis of a complete nitrifier reveals an oligotrophic lifestyle. *Nature*. 2017; 549:269–72.
13. Nowka B, Off S, Daims H, Spieck E. Improved isolation strategies allowed the phenotypic differentiation of two *Nitrospira* strains from widespread phylogenetic lineages. *FEMS Microbiol Ecol*. 2015;91(3):fiu031.
14. Ushiki N, Fujitani H, Aoi Y, Tsuneda S. Isolation of *Nitrospira* belonging to sublineage II from a wastewater treatment plant. *Microbes Environ*. 2013; 28(3):346–53.
15. van Kessel MA, Speth DR, Albertsen M, Nielsen PH, Op den Camp HJ, Kartal B, et al. Complete nitrification by a single microorganism. *Nature*. 2015; 528(7583):555–9.
16. Gruber-Dorninger C, Pester M, Kitzinger K, Savio DF, Loy A, Rattei T, et al. Functionally relevant diversity of closely related *Nitrospira* in activated sludge. *ISME J*. 2015;9(3):643–55.
17. Lückner S, Wagner M, Maixner F, Pelletier E, Koch H, Vacherie B, et al. A *Nitrospira* metagenome illuminates the physiology and evolution of globally important nitrite-oxidizing bacteria. *Proc Natl Acad Sci U S A*. 2010;107(30): 13479–84.
18. Ushiki N, Fujitani H, Shimada Y, Morohoshi T, Sekiguchi Y, Tsuneda S. Genomic Analysis of Two Phylogenetically Distinct *Nitrospira* Species Reveals Their Genomic Plasticity and Functional Diversity. *Front Microbiol*. 2018;8:2637.
19. Daims H, Lebedeva EV, Pjevac P, Han P, Herbold C, Albertsen M, et al. Complete nitrification by *Nitrospira* bacteria. *Nature*. 2015;528(7583):504–9.
20. Ehrich S, Behrens D, Lebedeva E, Ludwig W, Bock E. A new obligately chemolithoautotrophic, nitrite-oxidizing bacterium, *Nitrospira moscoviensis* sp. nov. and its phylogenetic relationship. *Arch Microbiol*. 1995;164(1):16–23.
21. Nowka B, Daims H, Spieck E. Comparison of oxidation kinetics of nitrite-oxidizing bacteria: nitrite availability as a key factor in niche differentiation. *Appl Environ Microbiol*. 2015;81(2):745–53.
22. Lückner S, Nowka B, Rattei T, Spieck E, Daims H. The genome of *Nitrospira gracilis* illuminates the metabolism and evolution of the major marine nitrite oxidizer. *Front Microbiol*. 2013;4:27.
23. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res*. 2015;25(7):1043–55.
24. Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A, et al. MicroScope - an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res*. 2013;41(D1):D636–47.
25. Vallenet D, Labarre L, Rouy Z, Barbe V, Bocs S, Cruveiller S, et al. MaGe: a microbial genome annotation system supported by synteny results. *Nucleic Acids Res*. 2006;34(1):53–65.
26. Wang Y, Coleman-Derr D, Chen G, Gu YQ. OrthoVenn: a web server for genome wide comparison and annotation of orthologous clusters across multiple species. *Nucleic Acids Res*. 2015;43(W1):W78–84.
27. Ushiki N, Jinno M, Fujitani H, Suenaga T, Terada A, Tsuneda S. Nitrite oxidation kinetics of two *Nitrospira* strains: the quest for competition and ecological niche differentiation. *J Biosci Bioeng*. 2017;123(5):581–9.
28. Maixner F, Noguera DR, Anneser B, Stoecker K, Wegl G, Wagner M, et al. Nitrite concentration influences the population structure of *Nitrospira*-like bacteria. *Environ Microbiol*. 2006;8(8):1487–95.
29. Spieck E, Ehrich S, Aamand J, Bock E. Isolation and immunocytochemical location of the nitrite-oxidizing system in *Nitrospira moscoviensis*. *Arch Microbiol*. 1998;169(3):225–30.
30. Starkenburg SR, Arp DJ, Bottomley PJ. Expression of a putative nitrite reductase and the reversible inhibition of nitrite-dependent respiration by nitric oxide in *Nitrobacter winogradskyi* Nb-255. *Environ Microbiol*. 2008; 10(11):3036–42.
31. Lawson CE, Lückner S. Complete ammonia oxidation: an important control on nitrification in engineered ecosystems? *Curr Opin Biotechnol*. 2018;50: 158–65.
32. Watson SW, Bock E, Valois FW, Waterbury JB, Schlosser U. *Nitrospira marina* gen. nov. sp. nov.: a chemolithotrophic nitrite-oxidizing bacterium. *Arch Microbiol*. 1986;144(1):1–7.
33. Krzywinski MI, Schein JE, Birol I, Connors J, Gascoyne R, Horsman D, et al. Circos: An information aesthetic for comparative genomics. *Genome Research*. 2009;19(9):1639–45. <https://doi.org/10.1101/gr.092759.109>.
34. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol*. 2008;26(5):541–7.
35. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci U S A*. 1990;87(12):4576–9.
36. Garrity GM, Holt JG, Spieck E, Bock E, Johnson DB, Spring S, et al. Phylum BVIII. Nitrospirae phy. nov. In: Boone DR, Castenholz RW, Garrity GM, editors. *Bergey's Manual® of Systematic Bacteriology: Volume One: The Archaea and the Deeply Branching and Phototrophic Bacteria*. New York: Springer New York; 2001. p. 451–64. https://doi.org/10.1007/978-0-387-21609-6_25.
37. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. *Nat Genet*. 2000;25:25.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

